



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Targeting glycolysis in proliferative kidney diseases

Ghazi, Susan ; Polesel, Marcello ; Hall, Andrew M

Abstract: Glycolytic activity is increased in proliferating cells, leading to the concept that glycolysis could be a therapeutic target in cystic diseases and kidney cancer. Preclinical studies using the glucose analog 2-deoxy-d-glucose have shown promise; however, inhibiting glycolysis in humans is unlikely to be without risks. While proximal tubules are predominantly aerobic, later segments are more glycolytic. Understanding exactly where and why glycolysis is important in the physiology of the distal nephron is thus crucial in predicting potential adverse effects of glycolysis inhibitors. Live imaging techniques could play an important role in the process of characterizing cellular metabolism in the functioning kidney. The goal of this review is to briefly summarize recent findings on targeting glycolysis in proliferative kidney diseases and to highlight the necessity for future research focusing on glycolysis in the healthy kidney.

DOI: <https://doi.org/10.1152/ajprenal.00460.2019>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-178491>

Journal Article

Accepted Version

Originally published at:

Ghazi, Susan; Polesel, Marcello; Hall, Andrew M (2019). Targeting glycolysis in proliferative kidney diseases. *American Journal of Physiology. Renal, Fluid and Electrolyte Physiology*, 317(6):F1531-F1535.

DOI: <https://doi.org/10.1152/ajprenal.00460.2019>

1 **Mini Review**

2 **Targeting Glycolysis in Proliferative Kidney Diseases**

3 **Susan Ghazi¹, Marcello Polesel¹, Andrew M Hall^{1,2}.**

4
5 ¹Institute of Anatomy, University of Zurich, Switzerland. ²Department of Nephrology, University
6 Hospital Zurich, Switzerland.

7
8 Address for correspondence:

9 Susan Ghazi, MSc
10 Institute of Anatomy
11 University of Zurich
12 Winterthurerstrasse 190
13 8057 Zurich
14 Switzerland
15 Email: susan.ghazi@anatomy.uzh.ch
16 Tel: +41 (0)44 635 53 52

17
18
19 Word count (excluding title page, figure legends and references): 2172

20
21 Running title: Glycolysis in kidney diseases

22
23 Keywords: Glycolysis, Proliferation, Metabolism, Kidney

24
25 One sentence summary: Glycolysis is a potential therapeutic target in proliferative kidney
26 diseases, but is also important for normal function in the distal nephron.

34 **Abstract**

35 Glycolytic activity is increased in proliferating cells, leading to the concept that glycolysis could
36 be a therapeutic target in cystic diseases and kidney cancer. Pre-clinical studies using the glucose
37 analogue 2-deoxy-D-glucose have shown promise; however, inhibiting glycolysis in humans is
38 unlikely to be without risks. While proximal tubules are predominantly aerobic, later segments
39 are more glycolytic. Understanding exactly where and why glycolysis is important in the
40 physiology of the distal nephron is thus crucial in predicting potential adverse effects of
41 glycolysis inhibitors. Live imaging techniques could play an important role in the process of
42 characterizing cellular metabolism in the functioning kidney. The goal of this mini-review is to
43 briefly summarize recent findings on targeting glycolysis in proliferative kidney diseases, and to
44 highlight the necessity for future research focusing on glycolysis in the healthy kidney.

45

Introduction

GLYCOLYSIS is the metabolic pathway that converts glucose into pyruvate, alongside this process ATP and NADH are generated. In the presence of oxygen, pyruvate is actively transported into the mitochondria and further metabolized in the tricarboxylic acid (TCA) cycle (17), which produces energy precursors for oxidative phosphorylation. In the absence of oxygen, pyruvate is converted into lactate, and glycolysis remains the main source of ATP production, so-called anaerobic glycolysis (32).

Oxygen availability in the kidney decreases gradually from the cortex to the medulla, and this is associated with a difference in metabolism between the cortical and medullary nephron segments (12). Proximal tubules in the cortex perform the bulk of fluid reabsorption and are known to be highly aerobic (54), since oxidative phosphorylation is the most efficient method to generate ATP required to drive solute transport. In contrast, distal nephron segments are much more glycolytic (15, 54). In the medulla this is probably partly explained by the lower oxygen tension, but it is also possible that glycolysis might be advantageous for some specialized functions in the cortical distal nephron segments.

Glycolysis is known to be upregulated in dividing cells in the normal development of the kidney (22), and similar findings have recently also been observed in proliferative kidney diseases. Glycolysis is therefore increasingly being investigated as a therapeutic target for cystic diseases and kidney cancer (6, 24, 42). However, given the importance of glycolysis in normal kidney function, inhibiting this process is not without risks. This mini-review will briefly summarize the experience to date of targeting glycolysis in proliferative kidney diseases, and will highlight the need for new research to better understand where and why glycolysis is important in the normal physiology of the distal nephron.

Glycolysis in proliferative kidney diseases

The “Warburg effect”

In highly proliferative cells, such as cancer cells, glycolysis is typically used as energy source even when sufficient levels of oxygen are available to perform oxidative phosphorylation. This phenomenon is called aerobic glycolysis or the “Warburg effect”, initially described by Otto Warburg almost one hundred years ago (23, 50). Although not as efficient for producing ATP, glycolysis is beneficial for proliferative cells, as their energy requirements are typically low and glucose can be utilized instead for anabolic processes necessary for new cell formation (21, 23). In the following sections, we will briefly highlight two important proliferative kidney diseases, where evidence suggests glycolysis is also upregulated.

Polycystic kidney disease

Autosomal dominant polycystic kidney disease (ADPKD) is a chronic, inherited disorder that is characterized by fluid filled cysts originating in a minority of nephrons, and is caused by mutations in polycystin 1 (*Pkd1*) or 2 (*Pkd2*) (10, 14). Cell proliferation is one of the major factors contributing to cyst progression, and activation of cell proliferation signalling pathways has repeatedly been identified in ADPKD (18, 20, 38, 41, 49). In the last decade numerous studies have reported evidence of upregulated glycolysis in the pathogenesis of ADPKD (7, 24, 31, 37, 38), and this subject has recently been comprehensively reviewed by Padovano *et al.* (31). Briefly, Rowe *et al.* were one of the first to describe an alteration in metabolism in ADPKD, and that mouse embryonic fibroblasts lacking *Pkd1* rely on glycolysis as an energy source, even under aerobic conditions (38). These findings were supported by a subsequent study showing an upregulation of the glycolytic enzymes hexokinase 1 and 2 (*Hk1*, *Hk2*) and lactate dehydrogenase A (*LDHa*) in a rat model of ADPKD (37). Further *in vivo* evidence for the Warburg effect came

from a *Pkd1* inducible knock-out model, where an increased glucose uptake and conversion to lactate was observed, by following the ¹³C-labelled forms of these molecules (7). It is important to note that not all studies have detected increased glycolysis in ADPKD (26, 51). However, it seems clear that substantial changes in metabolism occur in tubular cells during cyst development, and understanding the nature of these is likely to be crucial to fully elucidating the pathogenesis.

Kidney cancer

Consistent with the Warburg effect, glycolysis is also thought to be upregulated in clear cell renal cell carcinoma (ccRCC) (5, 33, 34, 48, 52). This is the most common type of cancer in the kidney, and is mainly caused by a mutation in the tumour suppressor gene *VHL* (29, 39). The degree of shift towards glycolysis seems to be grade dependent, and is most marked in high-grade disease (5). Upregulation of glycolysis is associated with a simultaneous decrease in expression of TCA cycle genes, but an increase in genes involved in glutamine transport and fatty acid synthesis (5). These findings are likely explained by studies performed in *VHL*-deficient ccRCC cells, which showed that the TCA acts in reverse (so-called reductive carboxylation), using glutamine to generate citrate, which can then be exported to the cytosol to increase the production of macromolecules needed for cell proliferation (13). Interestingly, there is now evidence that this phenomenon might also occur in ADPKD (35), suggesting overlap in the disease mechanisms.

Glycolysis as treatment target

Given that glycolysis seems to be a crucial process in cell proliferation, it follows that inhibiting glycolysis might be beneficial in treating diseases like ADPKD and ccRCC, and numerous studies are now investigating this possibility. Aerobic glycolysis can potentially be inhibited on

many levels (reviewed in (27, 31, 42, 44)), including: the uptake of glucose into the cell by glucose transporters; the breakdown of glucose to glucose-6-phosphate by hexokinase; the formation of pyruvate by pyruvate kinase; and the conversion of pyruvate to lactate by LDH. Furthermore, besides glycolysis, the increased uptake of glutamine can also be targeted to inhibit reductive carboxylation (13, 35). However, the strategy that has received most attention is direct inhibition of glycolysis with the glucose analogue 2-deoxy-D-glucose (2DG).

2-deoxy-D-glucose

2DG is taken up by cells and phosphorylated by hexokinase, but is not further catabolized in the glycolysis pathway (53). Although widely used in cancer biology (55), it has not yet been extensively tested in pre-clinical ccRCC models. However, a recent study using primary cell cultures from low- and high-grade ccRCC showed that 2DG significantly decreased lactate secretion and ATP levels only in the latter (2). This suggests that the usage of glycolysis inhibition as a therapy might need to be tailored to individual patients, depending on information acquired from tissue histology.

Meanwhile, 2DG has also been tested in various *in vitro* and *in vivo* studies of ADPKD. Administration of 2DG significantly retarded cyst progression in two different rodent models (7, 37). Furthermore, a low dose combinational treatment of 2DG with metformin (normally used for the treatment of type 2 diabetes mellitus) was shown to inhibit cell proliferation in human-derived cystic kidney epithelial cells (56). This combination treatment was also tested in a miniature pig model, and was shown to markedly reduce renal cyst formation and protect renal function (19).

Overall, early studies have suggested that 2DG has some promise as a therapy for proliferative kidney disease. However, glycolysis is a fundamental and widespread metabolic process in living

organisms, and glucose depletion in the blood (hypoglycaemia) is known to produce serious adverse effects (43), including in the nervous, immune and respiratory systems (9, 25). Thus, intentionally inhibiting glycolysis *in vivo* brings significant potential risks, and careful attention needs to be given to safety issues.

Safety issues with blocking glycolysis in vivo

The use of 2DG has been reported in phase I clinical trials of different types of tumours and generally it seems to be surprisingly well tolerated (36, 43, 45). Therefore, it should be possible to implement its use in clinical trials of ADPKD patients (reviewed in (24)). However, the safety profile of this compound, together with the appropriate dose, needs to be more carefully determined. While some animal studies report no degree of toxicity over prolonged periods of administration of 2DG (7, 30), others have described severe adverse cardiac effects (28, 46). Moreover, cardiac abnormalities, together with disorientation, tiredness, and excessive sweating have also been observed in the human clinical trials (36, 45).

Finding a treatment regime that produces sufficient blockade of glycolysis in proliferating cells to substantially retard disease progression, but without adversely altering metabolism in healthy tissues, represents a major logistical challenge. This concept is particularly pertinent to the kidney, where levels of glycolysis are normally quite high in the medulla (Figure 1A). Moreover, besides producing ATP, glucose is also important for other metabolic processes, such as the generation of NAD(P)H via the pentose phosphate pathway. Indeed, we recently found that administration of 2-DG acutely decreases mitochondrial and cytosolic NAD(P)H signals in proximal tubules (4).

Thus, while inhibiting glycolysis could be of benefit in retarding the progression of kidney cysts and tumours, side effects might arise, partly due to a loss of normal specialized functions in distal nephron segments. To understand the nature and magnitude of these effects, it will first be necessary to pin-point which cell types and transport processes are particularly dependent on glycolysis. This in turn will necessitate the development of new techniques to study cellular metabolism in intact functioning kidney tissue.

Glycolysis in the distal nephron segments

Within the distal nephron, previous studies have suggested that the collecting duct (CD) is the site with the highest glycolytic activity. CDs express more glycolytic enzymes than other segments (15), generate more lactate (1), and maintain ATP levels better when aerobic respiration is inhibited (47). Unlike other segments, CDs comprise of two very different cell types interspersed with each other, which have very distinct transport functions. Principal cells (PCs) are responsible for sodium and water homeostasis, whereas intercalated cells (ICs) maintain an acid-base balance (8). In parallel with transport functions, it is likely that PCs and ICs also display differences in metabolism. However, relatively little is known about this, due to the technical challenges of isolating cells in an appropriate state to perform standard metabolic assays. Thus, most studies have historically considered the CD as a single entity, and it therefore remains unclear which cell type is predominantly responsible for the overall glycolytic phenotype, and also which specific transport processes might be dependent on glycolysis.

Morphological studies such as electron microscopy reveal major differences in ultrastructure between cells in the CD. Of relevance to metabolism, the density of mitochondria is much higher in ICs than in PCs (3) (Figure 1B), perhaps hinting that the latter might be more glycolytic.

188 However, mitochondria have important functions other than generating ATP, so it does not
189 necessarily follow that ICs generate more ATP via oxidative phosphorylation.

190 The development of high-resolution live imaging techniques like multiphoton microscopy
191 provides a potential new avenue to study cellular metabolism in living animals (16, 40). Using
192 this approach, it has recently been described that cells in the rat CD display striking differences in
193 the uptake of mitochondrial dyes, hinting at the existence of major differences in metabolism
194 between PCs and ICs (11). We have also observed similar signal patterns in mice (Figure 1C),
195 and are currently utilizing multiphoton microscopy - in parallel with other imaging techniques -
196 to try to further understand the significance of these findings, and what they might tell us about
197 the nature of metabolism along the CD. We envisage that such an approach could finally shed
198 some light on the relationship between glycolysis and transport function in the distal nephron,
199 understanding of which could be crucial in anticipating adverse effects of glycolysis inhibitors.

200
201 **Conclusion**

202 Glycolysis is upregulated in proliferating cells and has thus been identified as a novel therapeutic
203 target in ADPKD and kidney cancer. Pre-clinical studies with blocking agents such as 2DG have
204 shown some promise, but inhibiting glucose metabolism in humans is unlikely to be without risk.
205 In particular, glycolytic activity is thought to be high in segments like the CD. It is therefore
206 crucial to further elucidate where and why glycolysis is important in the nephron, and live cell
207 imaging techniques could play an important role in this process.

208
209 **Grants**

210 A.M.H. is supported by The Swiss National Centre for Competence in Research (NCCR) Kidney
211 Control of Homeostasis, and by a project grant from the Swiss National Science Foundation.

212

213 **Disclosures**

214 No conflicts of interest, financial or otherwise, are declared by the authors.

215

216 **Author contributions**

217 S.G. and M.P. prepared the figures; S.G. drafted manuscript; A.M.H. edited and revised
218 manuscript; A.M.H. approved final version of manuscript.

219

220 **References**

- 221 1. **Bagnasco S, Good D, Balaban R, Burg M.** Lactate production in isolated segments of the
222 rat nephron. *Am J Physiol - Ren Fluid Electrolyte Physiol* 17, 1985.
- 223 2. **Bianchi C, Meregalli C, Bombelli S, Di Stefano V, Salerno F, Torsello B, De Marco S,**
224 **Bovo G, Cifola I, Mangano E, Battaglia C, Strada G, Lucarelli G, Weiss RH, Perego**
225 **RA.** The glucose and lipid metabolism reprogramming is gradedependent in clear cell
226 renal cell carcinoma primary cultures and is targetable to modulate cell viability and
227 proliferation. *Oncotarget* 8: 113502–113515, 2017.
- 228 3. **Brown D, Breton S.** Mitochondria-rich, proton-secreting epithelial cells. *J. Exp. Biol.* 199:
229 2345–2358, 1996.
- 230 4. **Bugarski M, Martins JR, Haenni D, Hall AM.** Multiphoton imaging reveals axial
231 differences in metabolic autofluorescence signals along the kidney proximal tubule. *Am J*
232 *Physiol - Ren Physiol* 315: F1613–F1625, 2018.
- 233 5. **Cancer Genome Atlas Research Network.** Comprehensive molecular characterization of
234 clear cell renal cell carcinoma. *Nature* 499: 43–49, 2013.
- 235 6. **Cargill K, Sims-Lucas S.** Metabolic requirements of the nephron. *Pediatr. Nephrol.*
236 (2018). doi: 10.1007/s00467-018-4157-2.
- 237 7. **Chiaravalli M, Rowe I, Mannella V, Quilici G, Canu T, Bianchi V, Gurgone A,**
238 **Antunes S, D’Adamo P, Esposito A, Musco G, Boletta A.** 2-Deoxy-D-glucose
239 ameliorates PKD progression. *J Am Soc Nephrol* 27: 1958–1969, 2016.
- 240 8. **Christensen EI, Wagner CA, Kaissling B.** Uriniferous tubule: Structural and functional
241 organization. *Compr Physiol* 2: 805–861, 2012.

- 242 9. **Davis IC, Ahmadizadeh I, Randell J, Younk L, Davis SN.** Understanding the impact of
243 hypoglycemia on the cardiovascular system. *Expert Rev. Endocrinol. Metab.* 12: 21–33,
244 2017.
- 245 10. **Douguet D, Patel A, Honoré E.** Structure and function of polycystins: insights into
246 polycystic kidney disease. *Nat. Rev. Nephrol.* 15: 412–422, 2019.
- 247 11. **Dunn KW, Sutton TA, Sandoval RM.** Live-Animal Imaging of Renal Function by
248 Multiphoton Microscopy. *Curr Protoc Cytom* 83: 12.9.1–12.9.25, 2018.
- 249 12. **Epstein FH.** Oxygen and renal metabolism. In: *Kidney International*. 1997, p. 381–385.
- 250 13. **Gameiro PA, Yang J, Metelo AM, Pérez-Carro R, Baker R, Wang Z, Arreola A,**
251 **Rathmell WK, Olumi A, López-Larrubia P, Stephanopoulos G, Iliopoulos O.** In vivo
252 HIF-mediated reductive carboxylation is regulated by citrate levels and sensitizes VHL-
253 deficient cells to glutamine deprivation. *Cell Metab* 17: 372–385, 2013.
- 254 14. **Grantham JJ.** Clinical Practice. Autosomal Dominant Polycystic Kidney Disease. *N Engl*
255 *J Med* 359: 1477–85, 2008.
- 256 15. **Guder WG, Ross BD.** Enzyme distribution along the nephron. *Kidney Int* 26: 101–111,
257 1984.
- 258 16. **Hall AM, Crawford C, Unwin RJ, Duchon MR, Peppiatt-Wildman CM.** Multiphoton
259 imaging of the functioning kidney. *J Am Soc Nephrol* 22: 1297–304, 2011.
- 260 17. **Herzig S, Raemy E, Montessuit S, Veuthey JL, Zamboni N, Westermann B, Kunji**
261 **ERS, Martinou JC.** Identification and functional expression of the mitochondrial
262 pyruvate carrier. *Science (80-)* 336: 93–96, 2012.
- 263 18. **Ibraghimov-Beskrovnaya O, Natoli TA.** mTOR signaling in polycystic kidney disease.
264 *Trends Mol. Med.* 17: 625–633, 2011.
- 265 19. **Lian X, Wu X, Li Z, Zhang Y, Song K, Cai G, Li Q, Lin S, Chen X, Bai XY.** The
266 combination of metformin and 2-deoxyglucose significantly inhibits cyst formation in
267 miniature pigs with polycystic kidney disease. *Br J Pharmacol* 176: 711–724, 2019.
- 268 20. **Lian X, Zhao J, Wu X, Zhang Y, Li Q, Lin S, Bai XY, Chen X.** The changes in glucose
269 metabolism and cell proliferation in the kidneys of polycystic kidney disease mini-pig
270 models. *Biochem Biophys Res Commun* 488: 374–381, 2017.
- 271 21. **Liberti M V., Locasale JW.** The Warburg Effect: How Does it Benefit Cancer Cells?
272 *Trends Biochem. Sci.* 41: 211–218, 2016.
- 273 22. **Liu J, Edgington-Giordano F, Dugas C, Abrams A, Katakam P, Satou R, Saifudeen**
274 **Z.** Regulation of nephron progenitor cell self-renewal by intermediary metabolism. *J Am*
275 *Soc Nephrol* 28: 3323–3335, 2017.

- 276 23. **Lunt SY, Vander Heiden MG.** Aerobic Glycolysis: Meeting the Metabolic Requirements
277 of Cell Proliferation. *Annu Rev Cell Dev Biol* 27: 441–464, 2011.
- 278 24. **Magistroni R, Boletta A.** Defective glycolysis and the use of 2-deoxy-d-glucose in
279 polycystic kidney disease: from animal models to humans. *J. Nephrol.* 30: 511–519, 2017.
- 280 25. **McNay EC, Cotero VE.** Mini-review: Impact of recurrent hypoglycemia on cognitive and
281 brain function. *Physiol Behav* 100: 234–238, 2010.
- 282 26. **Menezes LF, Lin CC, Zhou F, Germino GG.** Fatty Acid Oxidation is Impaired in An
283 Orthologous Mouse Model of Autosomal Dominant Polycystic Kidney Disease.
284 *EBioMedicine* 5: 183–192, 2016.
- 285 27. **Van Der Mijn JC, Panka DJ, Geissler AK, Verheul HM, Mier JW.** Novel drugs that
286 target the metabolic reprogramming in renal cell cancer. *Cancer Metab* 4, 2016.
- 287 28. **Minor RK, Smith DL, Sossong AM, Kaushik S, Poosala S, Spangler EL, Roth GS,**
288 **Lane M, Allison DB, de Cabo R, Ingram DK, Mattison JA.** Chronic ingestion of 2-
289 deoxy-d-glucose induces cardiac vacuolization and increases mortality in rats. *Toxicol*
290 *Appl Pharmacol* 243: 332–339, 2010.
- 291 29. **Nickerson ML, Jaeger E, Shi Y, Durocher JA, Mahurkar S, Zaridze D, Matveev V,**
292 **Janout V, Kollarova H, Bencko V, Navratilova M, Szeszenia-Dabrowska N, Mates D,**
293 **Mukeria A, Holcatova I, Schmidt LS, Toro JR, Karami S, Hung R, Gerard GF,**
294 **Linehan WM, Merino M, Zbar B, Boffetta P, Brennan P, Rothman N, Chow WH,**
295 **Waldman FM, Moore LE.** Improved identification of von Hippel-Lindau gene alterations
296 in clear cell renal tumors. *Clin Cancer Res* 14: 4726–4734, 2008.
- 297 30. **Ockuly JC, Gielissen JM, Levenick C V., Zeal C, Groble K, Munsey K, Sutula TP,**
298 **Stafstrom CE.** Behavioral, cognitive, and safety profile of 2-deoxy-2-glucose (2DG) in
299 adult rats. *Epilepsy Res* 101: 246–252, 2012.
- 300 31. **Padovano V, Podrini C, Boletta A, Caplan MJ.** Metabolism and mitochondria in
301 polycystic kidney disease research and therapy. *Nat. Rev. Nephrol.* 14: 678–687, 2018.
- 302 32. **Papandreou I, Cairns RA, Fontana L, Lim AL, Denko NC.** HIF-1 mediates adaptation
303 to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* 3:
304 187–197, 2006.
- 305 33. **Perroud B, Ishimaru T, Borowsky AD, Weiss RH.** Grade-dependent proteomics
306 characterization of kidney cancer. *Mol Cell Proteomics* 8: 971–985, 2009.
- 307 34. **Perroud B, Lee J, Valkova N, Dhirapong A, Lin PY, Fiehn O, Kültz D, Weiss RH.**
308 Pathway analysis of kidney cancer using proteomics and metabolic profiling. *Mol Cancer*
309 5, 2006.

- 310 35. **Podrini C, Rowe I, Pagliarini R, Costa ASH, Chiaravalli M, Di Meo I, Kim H,**
311 **Distefano G, Tiranti V, Qian F, di Bernardo D, Frezza C, Boletta A.** Dissection of
312 metabolic reprogramming in polycystic kidney disease reveals coordinated rewiring of
313 bioenergetic pathways. *Commun Biol* 1, 2018.
- 314 36. **Raez LE, Papadopoulos K, Ricart AD, Chiorean EG, Dipaola RS, Stein MN, Rocha**
315 **Lima CM, Schlesselman JJ, Tolba K, Langmuir VK, Kroll S, Jung DT, Kurtoglu M,**
316 **Rosenblatt J, Lampidis TJ.** A phase I dose-escalation trial of 2-deoxy-d-glucose alone or
317 combined with docetaxel in patients with advanced solid tumors. *Cancer Chemother*
318 *Pharmacol* 71: 523–530, 2013.
- 319 37. **Riwanto M, Kapoor S, Rodriguez D, Edenhofer I, Segerer S, Wüthrich RP.** Inhibition
320 of aerobic glycolysis attenuates disease progression in polycystic kidney disease. *PLoS*
321 *One* 11, 2016.
- 322 38. **Rowe I, Chiaravalli M, Mannella V, Ulisse V, Quilici G, Pema M, Song XW, Xu H,**
323 **Mari S, Qian F, Pei Y, Musco G, Boletta A.** Defective glucose metabolism in polycystic
324 kidney disease identifies a new therapeutic strategy. *Nat Med* 19: 488–493, 2013.
- 325 39. **Sato Y, Yoshizato T, Shiraishi Y, Maekawa S, Okuno Y, Kamura T, Shimamura T,**
326 **Sato-Otsubo A, Nagae G, Suzuki H, Nagata Y, Yoshida K, Kon A, Suzuki Y, Chiba**
327 **K, Tanaka H, Niida A, Fujimoto A, Tsunoda T, Morikawa T, Maeda D, Kume H,**
328 **Sugano S, Fukayama M, Aburatani H, Sanada M, Miyano S, Homma Y, Ogawa S.**
329 Integrated molecular analysis of clear-cell renal cell carcinoma. *Nat Genet* 45: 860–867,
330 2013.
- 331 40. **Schuh CD, Haenni D, Craigie E, Ziegler U, Weber B, Devuyst O, Hall AM.** Long
332 wavelength multiphoton excitation is advantageous for intravital kidney imaging. *Kidney*
333 *Int* 89: 712–719, 2016.
- 334 41. **Shillingford JM, Murcia NS, Larson CH, Low SH, Hedgepeth R, Brown N, Flask CA,**
335 **Novick AC, Goldfarb DA, Kramer-Zucker A, Walz G, Piontek KB, Germino GG,**
336 **Weimbs T.** The mTOR pathway is regulated by polycystin-1, and its inhibition reverses
337 renal cystogenesis in polycystic kidney disease. *Proc Natl Acad Sci U S A* 103: 5466–
338 5471, 2006.
- 339 42. **Shuch B, Linehan WM, Srinivasan R.** Aerobic glycolysis: A novel target in kidney
340 cancer. *Expert Rev. Anticancer Ther.* 13: 711–719, 2013.
- 341 43. **Singh D, Banerji AK, Dwarakanath BS, Tripathi RP, Gupta JP, Mathew TL,**
342 **Ravindranath T, Jain V.** Optimizing cancer radiotherapy with 2-deoxy-D-glucose: Dose
343 escalation studies in patients with glioblastoma multiforme. *Strahlentherapie und Onkol*
344 181: 507–514, 2005.
- 345 44. **Srinivasan R, Ricketts CJ, Sourbier C, Linehan WM.** New strategies in renal cell
346 carcinoma: Targeting the genetic and metabolic basis of disease. *Clin Cancer Res* 21: 10–
347 17, 2015.

45. **Stein M, Lin H, Jeyamohan C, Dvorzhinski D, Gounder M, Bray K, Eddy S, Goodin S, White E, Dipaola RS.** Targeting tumor metabolism with 2-deoxyglucose in patients with castrate-resistant prostate cancer and advanced malignancies. *Prostate* 70: 1388–1394, 2010.
46. **Terse PS, Joshi PS, Bordelon NR, Brys AM, Patton KM, Arndt TP, Sutula TP.** 2-Deoxy-d-Glucose (2-DG)-Induced Cardiac Toxicity in Rat: NT-proBNP and BNP as Potential Early Cardiac Safety Biomarkers. *Int J Toxicol* 35: 284–93, 2016.
47. **Uchida S, Endou H.** Substrate specificity to maintain cellular ATP along the mouse nephron. *Am J Physiol - Ren Fluid Electrolyte Physiol* 255, 1988.
48. **Unwin RD, Craven RA, Harnden P, Hanrahan S, Totty N, Knowles M, Eardley I, Selby PJ, Banks RE.** Proteomic changes in renal cancer and co-ordinate demonstration of both the glycolytic and mitochondrial aspects of the Warburg effect. In: *Proteomics*. 2003, p. 1620–1632.
49. **Wahl PR, Serra AL, Le Hir M, Molle KD, Hall MN, Wüthrich RP.** Inhibition of mTOR with sirolimus slows disease progression in Han:SPRD rats with autosomal dominant polycystic kidney disease (ADPKD). *Nephrol Dial Transplant* 21: 598–604, 2006.
50. **Warburg O, Wind F, Negelein E.** The metabolism of tumors in the body. *J Gen Physiol* 8: 519–530, 1927.
51. **Warner G, Hein KZ, Nin V, Edwards M, Chini CCS, Hopp K, Harris PC, Torres VE, Chini EN.** Food restriction ameliorates the development of polycystic kidney disease. *J Am Soc Nephrol* 27: 1437–1447, 2016.
52. **Wettersten HI, Aboud OA, Lara PN, Weiss RH.** Metabolic reprogramming in clear cell renal cell carcinoma. *Nat. Rev. Nephrol.* 13: 410–419, 2017.
53. **Wick AN, Drury DR, Nakada HI, Wolfe JB.** Localization of the primary metabolic block produced by 2-deoxyglucose. *J Biol Chem* 224: 963–969, 1957.
54. **Wirthensohn G, Guder WG.** Renal substrate metabolism. *Physiol. Rev.* 66: 469–497, 1986.
55. **Zhang D, Li J, Wang F, Hu J, Wang S, Sun Y.** 2-Deoxy-D-glucose targeting of glucose metabolism in cancer cells as a potential therapy. *Cancer Lett.* 355: 176–183, 2014.
56. **Zhao J, Ma Y, Zhang Y, Fu B, Wu X, Li Q, Cai G, Chen X, Bai X-Y.** Low-dose 2-deoxyglucose and metformin synergically inhibit proliferation of human polycystic kidney cells by modulating glucose metabolism. *Cell Death Discov* 5, 2019.

383 **Figure legends**

384 **Figure 1: Glycolysis in the kidney and heterogeneity of the collecting duct.** (A) The oxygen
385 gradient in the kidney decreases gradually from the cortex to the medulla, and glycolytic activity
386 is higher in the latter. (B) Electron microscopy of a cortical collecting duct (CD), showing that
387 intercalated cells (ICs) have a much higher mitochondrial density than neighboring principal cells
388 (PCs). (C) *In vivo* multiphoton imaging of a mouse kidney reveals a heterogeneous uptake along
389 the cortical CD of the voltage dependent mitochondrial dye Tetramethylrhodamine (TMRM)
390 (red). TMRM signal intensity provides a readout of mitochondrial energization within living
391 cells, which is dependent on respiratory chain activity. Scale bar = 5 μm (A) and 10 μm (C).
392 Schematic image was produced using scientific illustration toolkits from Motifolio.

393

Figure 1

